

**Duality–Symmetric Eleven–Dimensional Supergravity
and its Coupling to M–Branes**

Igor Bandos¹, Nathan Berkovits² and Dmitri Sorokin¹

¹ NSC, Kharkov Institute of Physics and Technology
Kharkov, 310108, Ukraine
e-mail: bandos,dsorokin@kipt.kharkov.ua

² Instituto de Física Teórica, Univ. Estadual Paulista
Rua Pamplona 145, São Paulo, SP 01405-900, BRASIL
e-mail: nberkovi@ift.unesp.br

The standard eleven-dimensional supergravity action depends on a three-form gauge field and does not allow direct coupling to five-branes. Using previously developed methods, we construct a covariant eleven-dimensional supergravity action depending on a three-form and six-form gauge field in a duality-symmetric manner. This action is coupled to both the M-theory two-brane and five-brane, and corresponding equations of motion are obtained. Consistent coupling relates $D = 11$ duality properties with self-duality properties of the M-5-brane. From this duality-symmetric formulation, one derives an action describing coupling of the M-branes to standard $D = 11$ supergravity.

1. Introduction

Eleven-dimensional supergravity has recently returned to popularity because of M-theory conjectures. The standard $D=11$ supergravity action is constructed using a graviton, gravitino and three-form gauge field $A^{(3)}$ [1]. Although it is easy to couple the standard $D = 11$ supergravity multiplet to supergravity solutions corresponding to “electric” 2-branes [2], it is difficult to couple it to supergravity solutions corresponding to “magnetic” 5-branes [3]. But as the analysis of the effective worldvolume action of the M-theory five-brane has shown [4] [5] [6] [7] [8], the M-five-brane is a dyonic object which carries both an “electric” and a “magnetic” charge. Thus, for coupling the M-5-brane it is desirable to have a $D = 11$ supergravity action which would contain $A^{(3)}$ as well as a six-form gauge field $A^{(6)}$ whose field strength is the Hodge dual of the four-form field strength. The importance of the six-form gauge field in eleven-dimensional supergravity was first recognized in [9] and [10], and was later related in [11] to central charges in the M-theory superalgebra.

In [12], actions for the bosonic sector of $D = 11$ supergravity with $A^{(3)}$ and $A^{(6)}$ and their coupling to a membrane and a five-brane have been studied in an approach where the duality relations are imposed as extra constraints at the level of equations of motion. This approach seems to be not completely satisfactory since any modification of these actions (such as self-coupling, coupling to other fields and sources, and quantum corrections) would require corresponding consistent modification of the duality constraints, which can be hard to guess if these constraints are not yielded by the action. For instance, the five-brane action produces a highly non-linear self-duality condition for a two-form worldvolume gauge field [13] [5] which reduces to one used in [12] only in the linear approximation.

The situation with coupling in $D = 11$ supergravity is similar to that of the four-dimensional Maxwell action which is easily coupled to electric sources but not magnetic sources [14] [15] [16]. The covariant coupling of Maxwell theory to electric and magnetic sources has recently been studied [17] [18] using two different approaches.

The first approach requires an infinite number of fields and generalizes the McClain-Wu-Yu-Wotzasek action for two-dimensional chiral bosons [19] [20] [21][22]. The second approach was developed in [23] [24] and uses a harmonic-like variable (constructed of a scalar field derivative) to make manifestly covariant duality-symmetric actions [25] [26], a dual form of which was first studied by Zwanziger [15] in application to Maxwell theory. In $D=4$, both of these approaches introduce a second vector gauge field whose on-shell field strength is the dual to the original Maxwell field strength.

In this paper, these two approaches will be used to construct duality-symmetric actions for $D = 11$ supergravity which produce the duality relation between $A^{(3)}$ and $A^{(6)}$ as a consequence of their equations of motion. Then, by generalizing methods developed in [14][16][18], we couple the supergravity action of the second approach to the membrane and the 5-brane of M-theory. We observe an interesting phenomenon of intertwining local D=11 and worldvolume symmetries which are responsible, respectively, for duality properties of $D = 11$ supergravity and self-duality properties of the M-5-brane. Upon solving for part of the duality constraints, one can reduce these actions to the standard Cremmer-Julia-Scherk $D = 11$ supergravity and obtain its consistent coupling to the M-branes, but cannot produce a version with the gauge field $A^{(6)}$ alone.

Section 2 reviews the standard D=11 supergravity action. Section 3 describes the McClain-Wu-Yu-Wotzasek form of this action which then is truncated to a form containing a single auxiliary scalar field. This version of the theory is discussed in section 4 and, in section 5, it is coupled to the M-branes. Section 6 contains some concluding remarks and section 7 is an appendix which discusses supersymmetry transformations of $A^{(6)}$.

2. Six-form gauge fields in D=11 supergravity

2.1. Review of D=11 supergravity

Our notation and conventions are close to [27]. We use the almost plus signature for the metric and underlined latin letters for the indices of $D = 11$ vectors. Not underlined latin indices will correspond to M-brane worldvolumes.

The standard eleven-dimensional supergravity action is [1]:

$$\begin{aligned} \mathcal{S}_{CJS} = & \int d^{11}x \left[\frac{1}{4} e R(\omega) - \frac{ie}{2} \bar{\Psi}_{\underline{m}} \Gamma^{\underline{mnp}} D_{\underline{n}} \left[\frac{1}{2} (\omega + \tilde{\omega}) \right] \Psi_{\underline{p}} \right. \\ & - \frac{1}{4!7!} \varepsilon^{\underline{m}_1 \dots \underline{m}_4 \underline{n}_1 \dots \underline{n}_7} (C^{(7)} + *C^{(4)})_{\underline{n}_1 \dots \underline{n}_7} (F^{(4)} - \frac{1}{2} C^{(4)})_{\underline{m}_1 \dots \underline{m}_4} \\ & \left. - \frac{e}{2 \cdot 4!} F^{(4)}_{\underline{m}_1 \dots \underline{m}_4} F^{(4)\underline{m}_1 \dots \underline{m}_4} \right] + \int \frac{1}{3} A^{(3)} \wedge F^{(4)} \wedge F^{(4)}, \end{aligned} \quad (2.1)$$

where $e_{\underline{m}}^{\underline{a}}$ is a vielbein describing coupling to gravity, $e = \det e_{\underline{m}}^{\underline{a}}$ (letters from the beginning of the alphabet denote flat tangent space indices) and $g_{\underline{mn}} = e_{\underline{m}}^{\underline{a}} e_{\underline{n}}^{\underline{a}}$ is a D=11 metric; $F^{(4)} = dA^{(3)}$ is the field strength of the three-form gauge field, $\Psi_{\underline{m}}^{\alpha}(x)$ is the gravitino

field ($\alpha=1,\dots,32$), $\omega_{\underline{mab}}$ is a spin connection with torsion and $D_{\underline{m}}(\omega)$ is a corresponding covariant derivative as defined in [1][27], $\tilde{\omega}_{\underline{mab}} = \omega_{\underline{mab}} + \frac{i}{4}\bar{\Psi}^{\underline{n}}\Gamma_{\underline{mabn}}\Psi^{\underline{p}}$,

$$C_{\underline{n}_1\dots\underline{n}_7}^{(7)} = \frac{e}{4\cdot 4!}\varepsilon_{\underline{m}_1\dots\underline{m}_4\underline{n}_1\dots\underline{n}_7}\bar{\Psi}_{\underline{p}}\Gamma^{\underline{p}\underline{m}_1\dots\underline{m}_4}\Psi_{\underline{q}} = -\frac{21}{2}\bar{\Psi}_{[\underline{n}_1}\Gamma_{\underline{n}_2\dots\underline{n}_5}\Psi_{\underline{n}_7]}, \quad \text{and}$$

$$C_{\underline{m}_1\underline{m}_2\underline{m}_3\underline{m}_4}^{(4)} = 3\bar{\Psi}_{[\underline{m}_1}\Gamma_{\underline{m}_2\underline{m}_3}\Psi_{\underline{m}_4]},$$

where the antisymmetric product $\Gamma^{(p)}$ of p gamma-matrices $dx^{\underline{m}}e_{\underline{m}}^{\underline{a}}\Gamma_{\underline{a}}$ satisfies the Hodge duality condition

$$\Gamma^{(p)} = -(-1)^{\frac{(11-p)(10-p)}{2}} * \Gamma^{(11-p)}.$$

This action is invariant under gauge transformations of $A^{(3)}$ ($\delta A^{(3)} = d\phi^{(2)}$) and under the supersymmetry transformations:

$$\delta e_{\underline{m}}^{\underline{a}} = -i\bar{\epsilon}\Gamma^{\underline{a}}\Psi_{\underline{m}}, \quad \delta A_{\underline{m}_1\underline{m}_2\underline{m}_3}^{(3)} = \frac{3}{2}\bar{\epsilon}\Gamma_{[\underline{m}_1\underline{m}_2}\Psi_{\underline{m}_3]}, \quad (2.2)$$

$$\delta\Psi_{\underline{m}} = D_{\underline{m}}(\tilde{\omega})\epsilon - \frac{i}{3!4!}(\Gamma_{\underline{m}\underline{n}_1\dots\underline{n}_4} - 8\Gamma_{\underline{n}_1\underline{n}_2\underline{n}_3}\underline{g}_{\underline{n}_4\underline{m}})\tilde{F}^{(4)\underline{n}_1\dots\underline{n}_4}\epsilon,$$

where

$$\tilde{F}^{(4)} = dA^{(3)} - C^{(4)} \equiv F^{(4)} - C^{(4)} \quad (2.3)$$

is a supercovariant field strength in the sense that its supersymmetry variation does not contain derivatives of the supersymmetry parameter $\epsilon^\alpha(x)$ [1].

It is easy to couple (2.1) to electric 2-branes by adding the term

$$\mu_E \int_{\mathcal{M}_3} dy^{m_1} \wedge dy^{m_2} \wedge dy^{m_3} A_{m_1 m_2 m_3}^{(3)}(x(y)) \quad (m = 0, 1, 2),$$

where y parametrizes the three-dimensional worldvolume $\mathcal{M}_3$ spanned by the 2-brane, $A(x(y))$ is the pullback onto $\mathcal{M}_3$ of the $D = 11$ gauge field form and μ_E is the electric charge (or membrane tension).

However, as mentioned in the introduction, it is not straightforward to couple to 5-branes using the fields in (2.1) since minimal coupling of the M-5-brane is described by the term [4][5]

$$\mu_M \int_{\mathcal{M}_6} dy^{m_1} \wedge \dots \wedge dy^{m_6} A_{m_1\dots m_6}^{(6)}(x(y)) \quad (m = 0, 1, \dots, 5)$$

where y now parametrizes the six-dimensional worldvolume $\mathcal{M}_6$ spanned by the 5-brane, $A^{(6)}(x(y))$ is the pullback onto $\mathcal{M}_6$ of a $D = 11$ six-form gauge field whose field strength is dual to $F^{(4)}$, and μ_M is the magnetic charge (or five-brane tension).

2.2. Six-form gauge field

So one needs to find an action for D=11 supergravity containing a six-form gauge field whose field strength is dual to $F^{(4)}$. The duality relation between $A^{(6)}$ and $A^{(3)}$ must be slightly more complicated than $F^{(7)} = *F^{(4)}$ since the equation of motion for $A^{(3)}$ which follows from (2.1) is not $d^*F^{(4)} = 0$, but is instead

$$d^*F^{(4)} = -F^{(4)} \wedge F^{(4)} + dC^{(7)} + d^*C^{(4)}.$$

Since the equations of motion for $A^{(3)}$ should imply Bianchi identities for the dual field-strength, the appropriate duality condition for the field strength of $A^{(6)}$ is

$$dA^{(6)} = *F^{(4)} + A^{(3)} \wedge F^{(4)} - C^{(7)} - *C^{(4)}. \quad (2.4)$$

As shown in the appendix, $A^{(6)}$ must transform as

$$\delta A_{\underline{m}_1 \dots \underline{m}_6}^{(6)} = 3\epsilon \Gamma_{[\underline{m}_1 \dots \underline{m}_5} \Psi_{\underline{m}_6]} + \delta A_{[\underline{m}_1 \underline{m}_2 \underline{m}_3}^{(3)} A_{\underline{m}_4 \underline{m}_5 \underline{m}_6]}^{(3)} \quad (2.5)$$

in order for (2.4) to be preserved on-shell by spacetime-supersymmetry transformations.

It will be convenient to define in addition to (2.3) the supercovariant field strength

$$\tilde{F}^{(7)} = dA^{(6)} - A^{(3)} \wedge dA^{(3)} + C^{(7)} = F^{(7)} + C^{(7)} \quad (2.6)$$

where $F^{(7)} \equiv dA^{(6)} - A^{(3)} \wedge dA^{(3)}$. Then the duality relation (2.4) takes the form

$$\tilde{F}^{(7)} = *\tilde{F}^{(4)}. \quad (2.7)$$

For later consideration, it is also convenient to introduce duality related “generalized” field strengths

$$\begin{aligned} \tilde{\mathcal{F}}^{(7)} &\equiv \tilde{F}^{(7)} - *\tilde{F}^{(4)}; & \tilde{\mathcal{F}}^{(4)} &\equiv \tilde{F}^{(4)} + *\tilde{F}^{(7)}; \\ \tilde{\mathcal{F}}^{(7)} &= -*\tilde{\mathcal{F}}^{(4)}, & \tilde{\mathcal{F}}^{(4)} &= *\tilde{\mathcal{F}}^{(7)}, \end{aligned} \quad (2.8)$$

which are zero when the duality relation (2.7) is satisfied.

An action which yields (2.7) as an equation of motion can be constructed using two methods. The first method will be discussed in the following section, and the second method will be described in section 4.

3. Duality-symmetric action with an infinite number of fields

One obvious way to obtain (2.7) as an equation of motion is to add the term

$$\int d^{11}x L_0^{(4)}(\tilde{F}^{(4)} + {}^*\tilde{F}^{(7)}) = \int d^{11}x L_0^{(4)}\tilde{\mathcal{F}}^{(4)}$$

to the action of (2.1) where $L_0^{(4)}$ is an unconstrained four-form which acts as a Lagrange multiplier. However, the equation of motion from varying $A^{(6)}$ will imply that $L_0^{(4)}$ describes a propagating field. To eliminate this undesired propagating field, one needs to add to the action the infinite sum $\int d^{11}x \sum_{I=0}^{\infty} L_I^{(4)} L_{I+1}^{(4)}$ where $L_I^{(4)}$ for $I = 0$ to ∞ is an infinite set of four-form auxiliary fields. So the complete action is

$$\mathcal{S}_{CJS} + \int d^{11}x (L_0^{(4)}\tilde{\mathcal{F}}^{(4)} + \sum_{I=0}^{\infty} L_I^{(4)} L_{I+1}^{(4)}). \quad (3.1)$$

As discussed in references [21],[22] and [17], variation of $L_{I+1}^{(4)}$ implies that $L_I^{(4)} = 0$ for each I , assuming that only a finite number of $L_I^{(4)}$'s are non-zero. In other words, the only solution to the equations of motion containing a finite number of non-zero fields is when the original $D = 11$ supergravity fields are on-shell and when $L_I^{(4)} = 0$ for all I . The condition that only a finite number of fields are non-vanishing can be understood as a discretized version of the asymptotic boundary condition $L^{(4)}(x) = 0$ as $x \rightarrow \infty$. [28]

The supersymmetry transformations which leave (3.1) invariant are easily found to be¹

$$\delta L_{2I}^{(4)} = 0, \quad (3.2)$$

$$\delta L_{2I+1 \underline{p}_1 \dots \underline{p}_4} = -M_{\underline{m}_1 \dots \underline{m}_4 \alpha}^q N_{\underline{q} \underline{p}_1 \dots \underline{p}_4}^\alpha L_{2I}^{\underline{m}_1 \dots \underline{m}_4} + M_{\underline{p}_1 \dots \underline{p}_4 \alpha}^q N_{\underline{q} \underline{m}_1 \dots \underline{m}_4}^\alpha L_{2I+2}^{\underline{m}_1 \dots \underline{m}_4},$$

$$\delta e_{\underline{m}}^a = -i\bar{\epsilon}\Gamma^a \Psi_{\underline{m}}, \quad \delta A_{\underline{m}_1 \underline{m}_2 \underline{m}_3}^{(3)} = \frac{3}{2}\bar{\epsilon}\Gamma_{[\underline{m}_1 \underline{m}_2} \Psi_{\underline{m}_3]},$$

$$\delta A_{\underline{m}_1 \dots \underline{m}_6}^{(6)} = 3\bar{\epsilon}\Gamma_{[\underline{m}_1 \dots \underline{m}_5} \Psi_{\underline{m}_6]} + \delta A_{\underline{m}_1 \underline{m}_2 \underline{m}_3}^{(3)} A_{\underline{m}_4 \underline{m}_5 \underline{m}_6}^{(3)},$$

$$\delta \Psi_{\underline{m}} = D_{\underline{m}}(\tilde{\omega})\epsilon - \frac{i}{3!4!}(\Gamma_{\underline{m} \underline{n}_1 \dots \underline{n}_4} - 8\Gamma_{\underline{n}_1 \underline{n}_2 \underline{n}_3} g_{\underline{n}_4 \underline{m}})(\tilde{F}^{(4)\underline{n}_1 \dots \underline{n}_4} - 2L_0^{(4)\underline{n}_1 \dots \underline{n}_4})\epsilon,$$

where

$$M_{\underline{m}_1 \dots \underline{m}_4 \alpha}^q = d\tilde{\mathcal{F}}_{\underline{m}_1 \dots \underline{m}_4}/d\psi_{\underline{q}}^\alpha$$

¹ We would like to thank Hitoshi Nishino for noticing a problem with the original version of these supersymmetry transformations.

and

$$N_{\underline{q}\underline{p}_1\cdots\underline{p}_4}^\alpha = d(\delta\psi_{\underline{q}}^\alpha)/dL_0^{\underline{p}_1\cdots\underline{p}_4}.$$

Note that the supersymmetry transformation of $L_I^{(4)}$ vanishes in the linearized approximation and is “local” in the sense that it only depends on $L_J^{(4)}$ for $I - 1 \leq J \leq I + 1$. These properties are also true for the infinite set of Ramond-Ramond fields in the superstring field theory action of [21] for the Type II superstring.

Although this method for introducing six-form gauge fields into the $D = 11$ supergravity action appears somewhat trivial, it is interesting to note that closed superstring field theory uses precisely this method for describing Ramond-Ramond fields and their coupling to electric and magnetic D-branes. Since the Type IIA superstring is conjectured to come from dimensional reduction of some eleven-dimensional M-theory containing supergravity, perhaps the closed superstring field theory action of [21] comes from dimensional reduction of some eleven-dimensional action similar to (3.1). Note, however, that this dimensional reduction can not be straightforward since the three-form gauge field of (2.1) reduces to both Ramond-Ramond and NS-NS gauge fields in ten dimensions.

We now turn to the second covariant approach to the description of duality-symmetric fields. As was shown in [24], this formulation can be considered as a consistent covariant truncation of the infinite series of the auxiliary fields in (3.1) by putting all L_I with $I > 0$ to zero and choosing $L_{0\underline{n}_1\cdots\underline{n}_4}^{(4)} = \frac{2}{(\partial a)^2} \partial_{\underline{m}} a \tilde{\mathcal{F}}_{\underline{m}[\underline{n}_1\underline{n}_2\underline{n}_3}^{(4)} \partial_{\underline{n}_4]} a$, where $a(x)$ is a scalar auxiliary field which appears in the model in a nonpolynomial way. In some sense, what we have done is that we have hidden the infinite series into this nonpolynomiality. This leads to one of the forms of the $D = 11$ supergravity action to be considered in the next section.

4. Covariant duality-symmetric formulation with the $a(x)$ -field

4.1. Actions for the dual gauge fields

As we have seen in the previous section, to construct a covariant D=11 supergravity action with both $A^{(3)}$ and $A^{(6)}$ fields, one should introduce auxiliary fields. In this section, we shall apply the covariant approach of [23][24] where covariance is gained by the use of the single auxiliary scalar field $a(x)$. It has been proven convenient to introduce a time-like “harmonic” unit vector

$$v_{\underline{m}}(x) \equiv \frac{\partial_{\underline{m}} a(x)}{\sqrt{-(\partial^L a \partial_L a)^2}}, \quad v^{\underline{m}} v_{\underline{m}} = -1, \quad (4.1)$$

by means of which $a(x)$ enters the action (though we could just as well use a space-like vector $\partial_{\underline{m}}a(x)/\sqrt{(\partial^{\underline{L}}a\partial_{\underline{L}}a)^2}$).

For simplicity, in this and the next subsection we put the gravitino field to zero. This is reflected in the absence of ‘tilde’ over the field strengths (2.3), (2.6) and (2.8), which now do not contain $C^{(4)}$ and $C^{(7)}$.

We present three equivalent forms of the action for the dual $A^{(3)}$ and $A^{(6)}$ field, which are obtained from each other by reordering corresponding terms, each of them being useful for different purposes:

$$S_A^1 = \int d^{11}x \, e \left[\frac{1}{2 \cdot 3!} v_{\underline{p}} \, {}^* F^{(7)}{}_{\underline{p} \underline{m}_1 \underline{m}_2 \underline{m}_3} \mathcal{F}^{(4)}_{\underline{m}_1 \underline{m}_2 \underline{m}_3 \underline{q}} v_{\underline{q}} \right. \\ \left. + \frac{1}{2 \cdot 6!} v_{\underline{p}} \, {}^* F^{(4)}{}_{\underline{p} \underline{m}_1 \dots \underline{m}_6} \mathcal{F}^{(7)}_{\underline{m}_1 \dots \underline{m}_6 \underline{q}} v_{\underline{q}} + \int \frac{1}{6} F^{(7)} \wedge F^{(4)} \right] \quad (4.2)$$

$$S_A^2 = \int d^{11}x \, e \left[\frac{-1}{4 \cdot 4!} F^{(4)}_{\underline{m}_1 \dots \underline{m}_4} F^{(4)}{}_{\underline{m}_1 \dots \underline{m}_4} - \frac{1}{4 \cdot 7!} F^{(7)}_{\underline{m}_1 \dots \underline{m}_7} F^{(7)}{}_{\underline{m}_1 \dots \underline{m}_7} \right. \\ \left. - \frac{1}{4 \cdot 3!} v_{\underline{p}} \mathcal{F}^{(4)}_{\underline{p} \underline{m}_1 \underline{m}_2 \underline{m}_3} \mathcal{F}^{(4)}{}_{\underline{q} \underline{m}_1 \underline{m}_2 \underline{m}_3} v_{\underline{q}} - \frac{1}{4 \cdot 6!} v_{\underline{p}} \mathcal{F}^{(7)}_{\underline{p} \underline{m}_1 \dots \underline{m}_6} \mathcal{F}^{(7)}{}_{\underline{q} \underline{m}_1 \dots \underline{m}_6} v_{\underline{q}} \right] \\ + \int \frac{1}{6} F^{(7)} \wedge F^{(4)}. \quad (4.3)$$

$$S_A^3 = - \int d^{11}x \, \frac{e}{2 \cdot 4!} F^{(4)}_{\underline{m}_1 \dots \underline{m}_4} F^{(4)}{}_{\underline{m}_1 \dots \underline{m}_4} + \int \frac{1}{3} A^{(3)} \wedge F^{(4)} \wedge F^{(4)} \\ - \int dx^{11} \frac{e}{2 \cdot 3!} v_{\underline{p}} \mathcal{F}^{(4)}_{\underline{p} \underline{m}_1 \underline{m}_2 \underline{m}_3} \mathcal{F}^{(4)}{}_{\underline{q} \underline{m}_1 \underline{m}_2 \underline{m}_3} v_{\underline{q}}. \quad (4.4)$$

The form (4.2) of the action is the most suitable for deriving the equations of motion of the gauge fields and $a(x)$, and getting the duality relations (2.7).

The forms (4.2) and (4.3) are manifestly duality-symmetric with respect to $F^{(7)}$ and $F^{(4)}$. (Note that, because of the definition (2.6), $F^{(7)} \wedge F^{(4)} = -A^{(3)} \wedge F^{(4)} \wedge F^{(4)}$ up to a total derivative). It is convenient to consider a combination of (4.2) and (4.3) when checking local bosonic symmetries of the model which we present in the next subsection.

Finally, the action in the form (4.4) is one which we have obtained by truncating the infinite field action of section 3. Though not manifestly duality-symmetric, (4.4) is very close to the $A^{(3)}$ field Lagrangian in the Cremmer–Julia–Scherk action (2.1). It differs from the latter by the last term, the only place where the field strength of $A^{(6)}$ is contained. This form is the most appropriate for verifying local supersymmetry of the complete D=11 supergravity action in the formulation considered.

4.2. Local bosonic symmetries of the action and equations of motion

The actions (4.2)–(4.4) are general coordinate invariant and possess the ordinary gauge symmetries

$$\delta A^{(3)} = d\phi^{(2)}, \quad \delta A^{(6)} = d\phi^{(6)} + \phi^{(2)} \wedge F^{(4)}, \quad (4.5)$$

as well as additional local symmetries whose presence ensures the duality relations between the gauge fields [25][26], space–time covariance, and an auxiliary nature of $a(x)$ [23][24]. The corresponding local transformations of the fields are

$$\delta A^{(3)} = da \wedge \varphi^{(2)}, \quad \delta A^{(6)} = da \wedge \varphi^{(6)} + da \wedge \varphi^{(2)} \wedge A^{(3)}; \quad (4.6)$$

$$\delta a = \phi(x), \quad \delta A^{(3)} = -\frac{\phi}{\sqrt{-(\partial a)^2}} i_v \mathcal{F}^{(4)}, \quad (4.7)$$

$$\delta_\phi A^{(6)} = \frac{\phi}{\sqrt{-(\partial a)^2}} i_v \mathcal{F}^{(7)} + \delta_\phi A^{(3)} \wedge A^{(3)},$$

where

$$i_v \mathcal{F}^{(4)} \equiv \frac{1}{3!} v^{\underline{p}} \mathcal{F}_{\underline{p} \underline{m}_1 \underline{m}_2 \underline{m}_3}^{(4)} dx^{\underline{m}_1} \wedge dx^{\underline{m}_2} \wedge dx^{\underline{m}_3}, \quad (4.8)$$

$$i_v \mathcal{F}^{(7)} \equiv \frac{1}{6!} v^{\underline{p}} \mathcal{F}_{\underline{p} \underline{m}_1 \dots \underline{m}_6}^{(7)} dx^{\underline{m}_1} \wedge \dots \wedge dx^{\underline{m}_6}$$

are the 3– and the 6–form obtained by contracting the 4– and the 7–form field strength with $v^{\underline{p}}$ (4.1).

For varying the action with respect to $A^{(3)}$, $A^{(6)}$ and $a(x)$, we should know the variations of the field strengths which are

$$\delta F^{(4)} = d(\delta A^{(3)}), \quad \delta F^{(7)} = d(\delta A^{(6)}) - d(\delta A^{(3)} \wedge A^{(3)}) - 2\delta A^{(3)} \wedge F^{(4)}. \quad (4.9)$$

Then a general variation of the action is

$$\begin{aligned} \delta S_A = & \int d^{11}x \left[\frac{1}{3!7!} \varepsilon^{\underline{m} \underline{n}_1 \underline{n}_2 \underline{n}_3 \underline{m}_1 \dots \underline{m}_7} v_{\underline{m}} (\mathcal{F}_{\underline{n}_1 \underline{n}_2 \underline{n}_3 \underline{p}}^{(4)} v^{\underline{p}}) \delta F_{\underline{m}_1 \dots \underline{m}_7}^{(7)} \right. \\ & \left. + \frac{1}{4!6!} \varepsilon^{\underline{m} \underline{n}_1 \dots \underline{n}_6 \underline{m}_1 \dots \underline{m}_4} v_{\underline{m}} (\mathcal{F}_{\underline{n}_1 \dots \underline{n}_6 \underline{p}}^{(7)} v^{\underline{p}}) \delta F_{\underline{m}_1 \dots \underline{m}_4}^{(4)} \right] + \delta_{a(x)} S_A, \end{aligned} \quad (4.10)$$

or in terms of differential forms (and up to a total derivative)

$$\delta S_A = - \int [v \wedge i_v \mathcal{F}^{(4)} \wedge \delta F^{(7)} - \delta A^{(3)} \wedge d(v \wedge i_v \mathcal{F}^{(7)})] + \delta_{a(x)} S_A$$

$$\begin{aligned}
&= \int [(\delta A^{(6)} - \delta A^{(3)} \wedge A^{(3)} - \frac{\delta a}{\sqrt{-(\partial a)^2}} i_v \mathcal{F}^{(7)}) \wedge d(v \wedge i_v \mathcal{F}^{(4)}) \\
&+ (\delta A^{(3)} + \frac{\delta a}{\sqrt{-(\partial a)^2}} i_v \mathcal{F}^{(4)}) \wedge (d(v \wedge i_v \mathcal{F}^{(7)}) + 2v \wedge i_v \mathcal{F}^{(4)} \wedge F^{(4)})].
\end{aligned} \tag{4.11}$$

From (4.11) we get the equations of motion of $A^{(3)}$ and $A^{(6)}$:

$$d(v \wedge i_v \mathcal{F}^{(4)}) = 0, \quad d(v \wedge i_v \mathcal{F}^{(7)}) + 2v \wedge i_v \mathcal{F}^{(4)} \wedge F^{(4)} = 0. \tag{4.12}$$

As usual in this sort of models [25][26][23][24], these equations reduce to the duality conditions $\mathcal{F}^{(4)} = 0 = \mathcal{F}^{(7)}$ (2.7) (with zero gravitino part) upon gauge fixing the symmetries under (4.6), and the equation of motion of $a(x)$ is not independent but is a consequence of (4.12), which reflects its auxiliary nature [23][24].

Substituting variations (4.5)–(4.7) into (4.11) one can easily check that they indeed form local symmetries of the action.

Using the $\varphi^{(2)}$ -transformations in (4.6), if we now gauge fix $i_v \mathcal{F}^{(4)} = 0$ (but do not use the second equation in (4.12) containing $i_v \mathcal{F}^{(7)}$, which is dynamical), and substitute this condition into (4.4) we get the standard Cremmer–Julia–Scherk action. This explains how duality–symmetric actions reduce to conventional ones [26][23]. Note, however, that we cannot eliminate $A^{(3)}$ and get an action only in terms of $A^{(6)}$, since $A^{(3)}$ enters the actions (4.2)–(4.4) directly (i.e. not only through its field strength as $A^{(6)}$ does). This is why a $D = 11$ supergravity action with $A^{(6)}$ alone has not been constructed [9][10].

4.3. Complete duality-symmetric action for $D=11$ supergravity

To obtain the $D=11$ supergravity action in a duality–symmetric form with the scalar auxiliary field, we should replace the part of the Cremmer–Julia–Scherk action containing $A^{(3)}$ with one of the Lagrangians (4.2)–(4.4) appropriately modified due to the presence of the gravitino terms. Note that the inclusion of gravitino terms in the full action must respect the local bosonic symmetries (4.6) and (4.7). The appropriate action is

$$\begin{aligned}
S = \int d^{11}x & \left[\frac{1}{4} e R(\omega) - \frac{ie}{2} \bar{\Psi}_{\underline{m}} \Gamma^{\underline{mnp}} D_{\underline{n}} \frac{1}{2} (\omega + \tilde{\omega}) \Psi_{\underline{p}} \right. \\
& \left. - \frac{1}{2 \cdot 4! 7!} \varepsilon^{\underline{m}_1 \dots \underline{m}_4 \underline{n}_1 \dots \underline{n}_7} (C^{(7)} + {}^* C^{(4)})_{\underline{n}_1 \dots \underline{n}_7} (F^{(4)} - \frac{1}{2} C^{(4)})_{\underline{m}_1 \dots \underline{m}_4} \right] + \tilde{S}_A.
\end{aligned} \tag{4.13}$$

In (4.13), $\tilde{S}_A$ is one of the duality-symmetric actions (4.2), (4.3), where $F^{(7)}$ is replaced with

$$F^{(7)} + C^{(7)} + {}^* C^{(4)} \tag{4.14}$$

everywhere except in $F^{(7)} \wedge F^{(4)}$. If $\tilde{S}_A$ is chosen in the form (4.4), then the coefficient in front of $(C^{(7)} + *C^{(4)})$ in (4.13) acquires an additional factor 2 (i.e. is the same as in (2.1)).

From the action (4.13), one gets the duality relation (2.4) (or (2.7)) between (4.14) and $F^{(4)}$.

Local supersymmetry transformations under which the action (4.13) is invariant are:

$$\begin{aligned} \delta a = 0, \quad \delta e_{\underline{m}}^a &= -i\bar{\epsilon}\Gamma^a\Psi_{\underline{m}}, \quad \delta A_{\underline{m}_1\underline{m}_2\underline{m}_3}^{(3)} = \frac{3}{2}\bar{\epsilon}\Gamma_{[\underline{m}_1\underline{m}_2}\Psi_{\underline{m}_3]}, \\ \delta A_{\underline{m}_1\dots\underline{m}_6}^{(6)} &= 3\bar{\epsilon}\Gamma_{[\underline{m}_1\dots\underline{m}_5}\Psi_{\underline{m}_6]} + \delta A_{\underline{m}_1\underline{m}_2\underline{m}_3}^{(3)} A_{\underline{m}_4\underline{m}_5\underline{m}_6}^{(3)}. \end{aligned} \quad (4.15)$$

The transformations of e_m^a and $A^{(3)}$ are the same as in the standard version while, for the gravitino, we have

$$\begin{aligned} \delta\Psi_{\underline{m}} &= D_{\underline{m}}(\tilde{\omega})\epsilon \\ -\frac{i}{3!4!}(\Gamma_{\underline{m}\underline{n}_1\dots\underline{n}_4} - 8\Gamma_{\underline{n}_1\underline{n}_2\underline{n}_3}g_{\underline{n}_4\underline{m}})(\tilde{F}^{(4)}_{\underline{n}_1\dots\underline{n}_4} - 4v^{[\underline{n}_1}\tilde{\mathcal{F}}^{(4)}_{\underline{n}_2\underline{n}_3\underline{n}_4]p}v_{\underline{p}})\epsilon. \end{aligned} \quad (4.16)$$

Eq. (4.16) reduces to the standard supersymmetry transformations when $i_v\tilde{\mathcal{F}}^{(4)}$ is put to zero. Note that

$$\tilde{F}^{(4)} + v \wedge i_v\tilde{\mathcal{F}}^{(4)} = - *(\tilde{F}^{(7)} + v \wedge i_v\tilde{\mathcal{F}}^{(7)}). \quad (4.17)$$

Using this duality property one can rewrite (4.16) in a duality-symmetric form

$$\begin{aligned} \delta\Psi_{\underline{m}} &= D_{\underline{m}}(\tilde{\omega})\epsilon \\ -\frac{i}{2\cdot3!4!}(\Gamma_{\underline{m}\underline{n}_1\dots\underline{n}_4} - 8\Gamma_{\underline{n}_1\underline{n}_2\underline{n}_3}g_{\underline{n}_4\underline{m}})(\tilde{F}^{(4)}_{\underline{n}_1\dots\underline{n}_4} - 4v^{[\underline{n}_1}\tilde{\mathcal{F}}^{(4)}_{\underline{n}_2\underline{n}_3\underline{n}_4]p}v_{\underline{p}})\epsilon \\ +\frac{i}{3!7!}(\Gamma_{\underline{m}\underline{n}_1\dots\underline{n}_7} - \frac{7}{2}\Gamma_{\underline{n}_1\dots\underline{n}_6}g_{\underline{n}_7\underline{m}})(\tilde{F}^{(7)}_{\underline{n}_1\dots\underline{n}_7} + 6v^{[\underline{n}_1}\tilde{\mathcal{F}}^{(7)}_{\underline{n}_2\dots\underline{n}_7]p}v_{\underline{p}})\epsilon. \end{aligned} \quad (4.18)$$

To check that the supergravity action (4.13) is indeed invariant under the supersymmetry transformations (4.15) and (4.16), it is convenient to use the form (4.4) of the dual gauge field action. Then all standard terms in the supersymmetry variation of (4.13) which do not contain $\tilde{\mathcal{F}}^{(4)}$ vanish, as was proved by Cremmer, Julia and Scherk [1], while the variation terms which contain $\tilde{\mathcal{F}}^{(4)}$ have a structure similar to the standard terms with $F^{(4)}$ and, hence, cancel as well.

We should note that the tensor (4.17) is invariant under the gauge transformations (4.5) and (4.6), but its variation with respect to (4.7) vanishes only on the mass shell

(4.12). This indicates that the supersymmetry variations (4.16) of the gravitino commute with the bosonic transformations (4.7) only up to equations of motion.

A reason why the supersymmetry transformations for gravitino acquire additional terms with $\tilde{\mathcal{F}}$ is that the auxiliary field $a(x)$ is assumed to be invariant under the supersymmetry transformations ².

This implies that the anticommutator of two supertransformations contains not only the bosonic translation generator $P_{\underline{m}}$ (as in the ordinary case) but also a generator $\mathcal{G}$ of the local transformation (4.7) with a value of the parameter such that it cancels the general coordinate transformation when acting on $a(x)$, the form of the anticommutator of supercharges being

$$\{Q_\alpha, Q_\beta\} = (\Gamma^{\underline{m}})_{\alpha\beta} (P_{\underline{m}} - (\partial_{\underline{m}} a) \mathcal{G}). \quad (4.19)$$

Note that the gravitino is invariant under (4.7) by definition, and its supersymmetry transformations (and their commutator with (4.7)) close only on the mass shell. Instead of modifying the supersymmetry transformations for the gravitino, one might try to find a suitable supersymmetry transformation of $a(x)$. But because $a(x)$ enters the action in a specific way, it seems problematic to find such a transformation. So we use the modified supersymmetry transformations for fermions as has always been done in the models of this kind [26][24][29] ³.

5. Coupling to the M-branes

The super-p-branes naturally couple to supergravity fields propagating in curved target *superspace* (i.e. to superfields), and the requirement of local *kappa*-symmetry of the super-p-branes puts these superfields on the mass shell. In other words, the classical super-p-branes propagate in a supergravity background which satisfies supergravity equations of motion without sources. Thus, coupling of the complete D=11 supergravity action to the super-M-brane worldvolume actions requires additional study. This problem does not arise in the bosonic case, and in what follows we shall consider coupling of the bosonic part of the duality-symmetric $D = 11$ supergravity action to the worldvolume actions of the membrane [31] and the five-brane [5].

² See [29] for a detailed discussion of the modified supersymmetry transformations in duality-symmetric models.

³ The only known exception is a model of supersymmetric chiral bosons in $d = 2$ for which a standard superfield formulation was constructed [30].

For this, we use an approach first proposed by Dirac [14] for describing electromagnetic interactions of monopoles (see [32] as a review), and further developed in [16][18]. The Dirac approach uses the fact that a charged object (a particle, a string etc.) couples locally (and often minimally) to the gauge field whose charge this object carries, and it couples (in general) nonlocally to the dual gauge field strength by means of a nonphysical Dirac string (or a p-brane) which stems from the charged object.

Applying this method to the membrane we can couple it to both the infinite field form and the nonpolynomial form of the duality-symmetric $D = 11$ supergravity action.

As to the five-brane, it carries a two-form gauge field with a self-dual field strength in its worldvolume, hence to construct an effective worldvolume action for the five-brane one should apply one of the approaches discussed above. So far, the covariant five-brane action has been constructed only in the $a(x)$ -field form [5]. We will see that this allows one to almost straightforwardly couple the five-brane to the actions (4.2) – (4.4), but not to (3.1). For coupling to the latter, one should probably construct an infinite field form of the five-brane action by generalizing relevant results of [33] on a duality-symmetric formulation of Dirac–Born–Infeld theory in $D = 4$.

In the following subsections we shall discuss features of “ $a(x)$ -field” coupling of $D = 11$ supergravity and the M-branes.

5.1. Membrane coupling

The worldvolume action for a membrane propagating in a curved $D = 11$ background and coupled to $A^{(3)}$ is [31]:

$$S_{\mathcal{M}_3} = - \int_{\mathcal{M}_3} d^3y \sqrt{-\det(g_{mn})} + \frac{1}{2} \int_{\mathcal{M}_3} A^{(3)} \quad m, n = 0, 1, 2 \quad (5.1)$$

where

$$g_{mn}(y) = \frac{\partial x^{\underline{p}}}{\partial y_m} g_{\underline{p}\underline{q}}(x(y)) \frac{\partial x^{\underline{q}}}{\partial y_n} \quad (5.2)$$

is an induced worldvolume metric and $A^{(3)}(x(y))$ is the pullback of $A^{(3)}(x)$ onto $\mathcal{M}_3$ (as in subsection 2.1). For simplicity, we have put the membrane tension to one.

We can extend (5.1) to an integral over $D = 11$ space-time by inserting a δ -function closed 8-form

$$*J^3 = \frac{1}{3!8!} dx^{\underline{m}_1} \wedge \dots \wedge dx^{\underline{m}_8} \epsilon_{\underline{m}_1 \dots \underline{m}_8 \underline{n}_1 \underline{n}_2 \underline{n}_3} \int_{\mathcal{M}_3} d\hat{x}^{\underline{n}_1} \wedge d\hat{x}^{\underline{n}_2} \wedge d\hat{x}^{\underline{n}_3} \delta(x - \hat{x}(y)) \quad (5.3)$$

with the support on $\mathcal{M}_3$, $J^{(3)\underline{mnp}} = \frac{1}{\sqrt{-\det(g_{mn})}} \int_{\mathcal{M}_3} d\hat{x}^{\underline{m}} \wedge d\hat{x}^{\underline{n}} \wedge d\hat{x}^{\underline{p}} \delta(x - \hat{x}(y))$ being the membrane current minimally coupled to $A^{(3)}(x)$.

To couple the membrane action (5.1) to the duality-symmetric actions (4.2)–(4.4) we have to take care of the local symmetries (4.6), (4.7). These are preserved if the field strength $F^{(7)} = dA^{(6)} - A^{(3)} \wedge F^{(4)}$ in (4.2)–(4.4) (except for the Chern–Simons terms) is extended to

$$\hat{F}^{(7)} = F^{(7)} - *G^{(4)}(x), \quad (5.4)$$

where $G^{(4)}$ is defined by the equation

$$d *G^{(4)} = *J^{(3)}. \quad (5.5)$$

A solution to (5.5) is

$$G^{(4)\underline{m}_1 \dots \underline{m}_4}(x) = \frac{1}{\sqrt{-\det(g_{mn})}} \int_{\mathcal{M}_4} d\hat{x}^{\underline{m}_1} \wedge \dots \wedge d\hat{x}^{\underline{m}_4} \delta(x - \hat{x}(z)), \quad (5.6)$$

where the integration is performed over a four-dimensional surface $\mathcal{M}_4$ parametrized by z , whose boundary is the membrane worldvolume $\mathcal{M}_3 = \partial\mathcal{M}_4$. This is the generalization of the Dirac string [14] to a Dirac three-brane stemmed from the membrane, by means of which the latter couples to the dual gauge field strength $F^{(7)}$.

The complete action describing membrane coupling is therefore

$$S = \int d^{11}x \frac{1}{4} e R(\omega) + \hat{S}_A + S_{\mathcal{M}_3}, \quad (5.7)$$

where $\hat{S}_A$ is either (4.2) or (4.3) with $\hat{F}^{(7)}$ (5.4) instead of $F^{(7)}$ everywhere except of the Chern–Simons term $F^{(7)} \wedge F^{(4)}$. If $\hat{S}_A$ is in the form (4.4), the minimal coupling term in (5.1) doubles.

The equations of motion of $A^{(3)}$ and $A^{(6)}$ one gets from the variation of (5.7) reduce to the duality conditions

$$\hat{F}^{(7)} = F^{(7)} - *G^{(4)} = *F^{(4)}, \quad *\hat{F}^{(7)} = -F^{(4)} \quad (5.8)$$

whose Bianchi identities are the $D = 11$ gauge field equations with the membrane source

$$d*F^{(4)} + F^{(4)} \wedge F^{(4)} = *J^{(3)}, \quad d*\hat{F}^{(7)} = 0. \quad (5.9)$$

From varying the metric, one gets the Einstein equations:

$$R_{\underline{mn}} - \frac{1}{2}g_{\underline{mn}}R = \frac{1}{3}(F_{\underline{m}\underline{p}_1\underline{p}_2\underline{p}_3}^{(4)}F_{\underline{n}}^{(4)\underline{p}_1\underline{p}_2\underline{p}_3} - \frac{1}{8}g_{\underline{mn}}F_{\underline{p}_1\underline{p}_2\underline{p}_3\underline{p}_4}^{(4)}F^{(4)\underline{p}_1\underline{p}_2\underline{p}_3\underline{p}_4}) + T_{\underline{mn}}^2, \quad (5.10)$$

where the energy-momentum tensor of $A^{(3)}$ and $A^{(6)}$ reduces to that of $A^{(3)}$ after taking into account the duality relation (5.8), and the last term on the right hand side of (5.10) is the membrane energy-momentum tensor $T_{\underline{mn}}^2 = \sqrt{-\det g_{pq}}g^{mn}\partial_m x^{\underline{m}}\partial_n x^{\underline{n}}$. From varying the membrane coordinate, one gets the equation of motion of the membrane which is

$$\begin{aligned} \frac{\partial}{\partial y^{\underline{m}}}(\sqrt{-\det g_{pq}}g^{mn}(y)\partial_n x^{\underline{r}}g_{\underline{r}\underline{q}}) - \frac{1}{2}g^{mn}\partial_m x^{\underline{m}}\partial_n x^{\underline{n}}\frac{\partial}{\partial x^{\underline{q}}}g_{\underline{mn}}(x) = \\ = \frac{1}{2 \cdot 3!}\epsilon^{mnp}\partial_m x^{\underline{m}}\partial_n x^{\underline{n}}\partial_p x^{\underline{p}}F_{\underline{mnpq}}^{(4)}. \end{aligned} \quad (5.11)$$

As expected, we have derived the standard equations of motion for the bosonic fields of $D = 11$ supergravity with the membrane as a source.

5.2. Five-brane coupling

The worldvolume action for the M-theory five-brane propagating in a curved $D = 11$ background and coupled to $A^{(3)}$ and $A^{(6)}$ is [5][6][7]:

$$S_{\mathcal{M}_6} = \int_{\mathcal{M}_6} d^6 y \left[-\sqrt{-\det(g_{mn} + i\tilde{H}_{mn})} + \sqrt{-g}\frac{1}{4}v_l {}^*H^{lmn}H_{mnp}v^p \right] \quad (5.12)$$

$$-\frac{1}{2}\int_{\mathcal{M}_6} \left[A^{(6)} + H^{(3)} \wedge A^{(3)} \right], \quad m, n = 0, 1, \dots, 5,$$

where $g_{mn}(y)$ is now the induced metric of the five-brane worldvolume $\mathcal{M}_6$;

$$H^{(3)} = dB^{(2)} - A^{(3)} = \frac{1}{3!}dy^m \wedge dy^n \wedge dy^l H_{lmn}(y), \quad (5.13)$$

$$H_{lmn}(y) = 3(\partial_l B_{mn} + \partial_m B_{nl} + \partial_n B_{lm}) - A_{lmn}^{(3)}(x(y)) \quad (5.14)$$

is the field strength of the worldvolume gauge field $B_{mn}(y)$ which satisfies the generalized self-duality condition on the mass shell: [13],

$$\tilde{H}_{mn} \equiv v^l(y) {}^*H_{lmn} = (i_v {}^*H^{(3)})_{mn}, \quad {}^*H^{mnl} = \frac{1}{3!\sqrt{-g}}\varepsilon^{mnlpqr}H_{pqr}; \quad (5.15)$$

and

$$v_m(y) \equiv \frac{\partial_m a(y)}{\sqrt{-(\partial^l a \partial_l a)^2}}, \quad v^m v_m = -1, \quad (5.16)$$

where $a(y) \equiv a(x(y))$ is the pullback onto $\mathcal{M}_6$ of the auxiliary scalar field $a(x)$ which we used in section 4 to construct the covariant duality-symmetric $D = 11$ supergravity action. In the 5-brane action, the field $a(y)$ plays the same role [5] as in the $D = 11$ action. We should stress that *a priori* one could try to use an independent worldvolume scalar field to ensure the covariance of the five-brane action but, as we shall see below, it turns out crucial for the consistent coupling of this action to the duality-symmetric $D = 11$ action (4.2) – (4.4) that the auxiliary worldvolume field is the pullback of $a(x)$.

The action (5.12) is manifestly invariant under general coordinate transformations of $\mathcal{M}_6$, and under the following local transformations which transform $B^{(2)}$ and $a(y)$:

$$\delta B^{(2)} = d\phi^{(1)}(y) + \phi^{(2)}(x(y)) + da \wedge \varphi^{(1)}(y) - \frac{\varphi(x(y))}{\sqrt{-(\partial a)^2}} \mathcal{H}^{(2)}, \quad \delta a = \varphi(x(y)), \quad (5.17)$$

where $\phi^{(1)}(y)$ is the one-form parameter of the standard gauge transformations of $B^{(2)}$, $\phi^{(2)}(x(y))$ is the pullback of the $D = 11$ gauge transformations (4.5) of $A^{(3)}$ which ensures that the field strength $H^{(3)}$ (5.13) is invariant under these transformations, $\varphi^{(1)}(y)$ parametrizes a worldvolume analog of the transformations (4.6) and implies an on shell self-duality of $H^{(3)}$, $\varphi(x(y))$ is the pullback of the corresponding $D = 11$ scalar parameter of the local transformations (4.7), and

$$\mathcal{H}^{(2)} \equiv i_v H^{(3)} - \frac{dy^n \wedge dy^m}{\sqrt{-\det g_{pq}}} \frac{\delta \sqrt{\det(g_{pq} + i\tilde{H}_{pq})}}{\delta \tilde{H}_{mn}}. \quad (5.18)$$

From (4.6), (4.7) and (5.17), we see that $D = 11$ and $\mathcal{M}_6$ local symmetries responsible for the duality properties of eleven-dimensional supergravity and of the M-five-brane are intrinsically related to each other.

The generalized non-linear self-duality condition on $H^{(3)}$,

$$\mathcal{H}^{(2)} = 0, \quad (5.19)$$

is a consequence of the equation of motion of $B^{(2)}$ which follows from (5.12) (see [13], [5] for details).

We can extend (5.12) to an integral over $D = 11$ space-time by inserting into it a δ -function closed 5-form

$$*J^6 = \frac{1}{5!6!} dx^{\underline{m}_1} \wedge \dots \wedge dx^{\underline{m}_5} \epsilon_{\underline{m}_1 \dots \underline{m}_5 \underline{n}_1 \dots \underline{n}_6} \int_{\mathcal{M}_6} d\hat{x}^{\underline{n}_1} \wedge \dots \wedge d\hat{x}^{\underline{n}_6} \delta(x - \hat{x}(y)) \quad (5.20)$$

with the support on $\mathcal{M}_6$, $J^{(6)\underline{q}_1 \dots \underline{q}_6} = \frac{1}{\sqrt{-\det(g_{mn})}} \int_{\mathcal{M}_6} d\hat{x}^{\underline{q}_1} \wedge \dots \wedge d\hat{x}^{\underline{q}_6} \delta(x - \hat{x}(y))$ being the five-brane current minimally coupled to $A^{(6)}(x)$.

To couple the five-brane action (5.1) to the duality-symmetric actions (4.2), (4.3) we have to take care not only of the local symmetries (4.6), (4.7), but also of (5.17). Then the action describing consistent coupling is

$$S = \int d^{11}x \frac{1}{4} e R(\omega) + \hat{S}_A + S_{\mathcal{M}_6} - \frac{1}{2} \int H^{(3)} \wedge \hat{F}^{(4)} \wedge *G^{(7)}. \quad (5.21)$$

In (5.21), $\hat{S}_A$ is either (4.2) or (4.3) where everywhere, except in the Chern-Simons term, $F^{(4)}$ and $F^{(7)}$ are replaced with

$$\hat{F}^{(4)} = F^{(4)} - *G^{(7)}, \quad \hat{F}^{(7)} = F^{(7)} - H^{(3)} \wedge *G^{(7)}, \quad (5.22)$$

and the seven-form $G^{(7)}(x)$ is defined by the equation

$$d *G^{(7)} = *J^{(6)}. \quad (5.23)$$

A solution to (5.5) is

$$G^{(7)\underline{m}_1 \dots \underline{m}_7}(x) = \frac{1}{\sqrt{-\det(g_{mn})}} \int_{\mathcal{M}_7} d\hat{x}^{\underline{m}_1} \wedge \dots \wedge d\hat{x}^{\underline{m}_7} \delta(x - \hat{x}(z)), \quad (5.24)$$

where the integration is performed over a seven-dimensional surface $\mathcal{M}_7$ parametrized by z , whose boundary is the five-brane worldvolume $\mathcal{M}_6 = \partial\mathcal{M}_7$. This is the generalization of the Dirac string [14] to a Dirac six-brane stemmed from the five-brane, by means of which the latter couples to the dual gauge field potentials $F^{(4)}$ and $F^{(7)}$.

To check that the action (5.21) contains the symmetries (4.5) – (4.7) and (5.17) (where $\delta A^{(6)}$ acquires the additional term $(da \wedge \varphi^{(1)} - \frac{\delta a}{\sqrt{-(\partial a)^2}} \mathcal{H}^{(2)}) \wedge *G^{(7)}$), we present the differential form of the variation of (5.21) with respect to $A^{(3)}$, $A^{(6)}$, $B^{(2)}$ and a :

$$\begin{aligned} \delta S = \int [& (\delta A^{(6)} - \delta A^{(3)} \wedge A^{(3)} - \delta B^{(2)} \wedge *G^{(7)} - \frac{\delta a}{\sqrt{-(\partial a)^2}} i_v \hat{\mathcal{F}}^{(7)}) \wedge d(v \wedge i_v \hat{\mathcal{F}}^{(4)}) \\ & + (\delta A^{(3)} + \frac{\delta a}{\sqrt{-(\partial a)^2}} i_v \hat{\mathcal{F}}^{(4)}) \wedge (d(v \wedge i_v \hat{\mathcal{F}}^{(7)}) + 2v \wedge i_v \hat{\mathcal{F}}^{(4)} \wedge F^{(4)} \\ & \quad - v \wedge \mathcal{H}^{(2)} \wedge *J^6 - v \wedge i_v \hat{\mathcal{F}}^{(4)} \wedge *G^{(7)}) \\ & - (\delta B^{(2)} + \frac{\delta a}{\sqrt{-(\partial a)^2}} \mathcal{H}^{(2)}) \wedge (d(v \wedge \mathcal{H}^{(2)}) + v \wedge i_v \hat{\mathcal{F}}^{(4)}) \wedge *J^{(6)}]. \end{aligned} \quad (5.25)$$

The equations of motion of $A^{(3)}$ and $A^{(6)}$ one gets from the variation of (5.21) reduce to the duality conditions

$$\hat{F}^{(7)} = * \hat{F}^{(4)}, \quad * \hat{F}^{(7)} = -\hat{F}^{(4)}$$

whose Bianchi identities are the $D = 11$ gauge field equations with the five-brane source

$$d* \hat{F}^{(4)} + F^{(4)} \wedge \hat{F}^{(4)} = H^{(3)} \wedge * J^{(6)}, \quad (5.26)$$

$$d* \hat{F}^{(7)} = * J^{(6)}.$$

The $D = 11$ Einstein equations take the form:

$$R_{\underline{mn}} - \frac{1}{2} g_{\underline{mn}} R = \frac{1}{3} (\hat{F}_{\underline{m} \underline{p}_1 \underline{p}_2 \underline{p}_3}^{(4)} \hat{F}_{\underline{n}}^{(4) \underline{p}_1 \underline{p}_2 \underline{p}_3} - \frac{1}{8} g_{\underline{mn}} \hat{F}_{\underline{p}_1 \underline{p}_2 \underline{p}_3 \underline{p}_4}^{(4)} \hat{F}^{(4) \underline{p}_1 \underline{p}_2 \underline{p}_3 \underline{p}_4}) + T_{\underline{mn}}^5, \quad (5.27)$$

where $T^{5 \underline{mn}} = \sqrt{-\det g_{pq}} T^{mn} \partial_m x^{\underline{m}} \partial_n x^{\underline{n}}$ is the energy-momentum tensor of the 5-brane. An explicit form of a ‘formal’ $d = 6$ energy-momentum tensor T^{mn} and equations of motion of the 5-brane coordinate $x^{\underline{m}}(y)$ can be derived from [34] [35]. We should note that the equation of motion (5.26) differs from an analogous equation considered in [12] by a term $*G^{(7)} \wedge *G^{(7)}$ which is absent in our version. Thus, eqs. (5.26) are selfconsistent without the extra assumption of [12] that $d *G^{(7)} \wedge *G^{(7)} = 0$. And, as we have already mentioned, the equation of motion of $B^{(2)}$ produces the nonlinear self-duality condition (5.19), while [12] used its linearized approximation (i.e. $H^{(3)} = *H^{(3)}$) which was imposed by hand.

We can reduce the action (5.21) to an action which describes coupling of the M-five-brane to the bosonic sector of the standard D=11 supergravity action. For this, we should first rewrite (5.21) in such a way that it will contain $\hat{S}_A$ in the form of (4.4) with hatted $\hat{F}^{(4)}$ and $\hat{F}^{(7)}$ (5.22) everywhere except in the Chern-Simons term. This changes the form of five-brane coupling, and the action takes the form

$$\begin{aligned} S = & \int d^{11}x \left[\frac{1}{4} e R - \frac{e}{2 \cdot 4!} \hat{F}_{\underline{m}_1 \dots \underline{m}_4}^{(4)} \hat{F}^{(4) \underline{m}_1 \dots \underline{m}_4} \right] + \int \frac{1}{3} A^{(3)} \wedge dA^{(3)} \wedge dA^{(3)} \\ & - \int d^{11}x \frac{e}{2 \cdot 3!} v^{\underline{p}} \hat{\mathcal{F}}_{\underline{p} \underline{m}_1 \underline{m}_2 \underline{m}_3}^{(4)} \hat{\mathcal{F}}^{(4) \underline{q} \underline{m}_1 \underline{m}_2 \underline{m}_3} v_{\underline{q}} \\ & - \int_{\mathcal{M}_6} d^6y \left[\sqrt{-\det(g_{mn} + i \tilde{H}_{mn})} - \frac{\sqrt{-g}}{4} v_l *H^{lmn} H_{mnp} v^p \right] \\ & - \frac{1}{2} \int_{\mathcal{M}_6} dB^{(2)} \wedge A^{(3)} + \frac{1}{2} \int A^{(3)} \wedge dA^{(3)} \wedge *G^{(7)}. \end{aligned} \quad (5.28)$$

We see that the term of (5.12) which described minimal coupling of the five-brane to $A^{(6)}$ is replaced in (5.28) with a nonminimal term $A^{(3)} \wedge dA^{(3)} \wedge *G^{(7)}$, and $A^{(6)}$ remains only inside of $\hat{\mathcal{F}}^{(4)}$. Now, imposing the gauge fixing condition $i_v \hat{\mathcal{F}}^{(4)} = 0$ (as in the free supergravity case of subsection 4.2) we can eliminate the term with $A^{(6)}$ from (5.28), and the remaining action describes consistent nonminimal coupling of the M-five-brane to the standard D=11 supergravity. This action reduces to a corresponding version proposed in [12] when the 5-brane action is replaced with a quadratic term for $H^{(3)}$ ⁴.

Finally, coupling to D=11 supergravity of both a membrane and a five-brane, with the membrane ending on the five-brane, is described by the combination of actions (5.7) and (5.21) (or (5.28)) to which (as in [12]) one must add the membrane boundary term

$$\int_{\partial\mathcal{M}_3} B^{(2)} \equiv \int_{\mathcal{M}_6} B^{(2)} \wedge *j^{(2)} \quad (5.29)$$

and extend the five-brane field strength (5.13) to $\hat{H}^{(3)} = H^{(3)} - *G^{(3)}$, where $d*G^{(3)} = *j^{(2)}$ and $j^{(2)mn}(y^6) = \frac{1}{\sqrt{-\det g_{pq}(y^6)}} \int_{\mathcal{M}_2} d\hat{y}^{6m} \wedge d\hat{y}^{6n} \delta(y^6 - \hat{y}^6(y^2))$ with y^6 and y^2 being, respectively, coordinates of $\mathcal{M}_6$ and $\mathcal{M}_2 = \partial\mathcal{M}_3$.

6. Conclusion

We have constructed a manifestly duality-symmetric formulation of $D = 11$ supergravity with the gauge fields $A^{(3)}$ and $A^{(6)}$, coupled its bosonic sector to the two-brane and the five-brane of M-theory, obtained corresponding equations of motion of the system, and shown how the M-five-brane couples to Cremmer-Julia-Scherk supergravity. As further generalization, one can consider coupling of $D = 11$ supergravity to intersecting M-branes.

We have found that consistent coupling of the five-brane worldvolume effective action to the supergravity action requires local symmetries responsible for duality properties of the two actions to be related to each other through the same auxiliary field.

The action for $D = 11$ supergravity coupled to the M-branes might be useful for a development of results [36] [12][37] in studying anomalies in M-theory.

Methods developed in this paper can possibly be applied to the study of coupling of a self-dual IIB $D = 10$ supergravity action [38] to D-branes, of their anomalies [39]

⁴ In the original version of this paper, it was mistakenly claimed that the $A^{(3)} \wedge dA^{(3)} \wedge *G^{(7)}$ term is absent from the action in [12].

and vacuum energy [40], in particular, in the case of a self-dual D-3-brane [41] which admits a manifest duality-symmetric description [33][42], as well as to the consideration of analogous problems in Type IIA superstring theory.

We should note that in this paper we dealt with classical systems. As has been stressed by Witten [36], one should expect to find a problem when quantizing any action for a self-dual system. For example, for a $d = 2$ chiral scalar, the path integral can not be defined in a modular-invariant manner, implying that the amplitude must depend on more than just the metric of the two-dimensional surface. In the first approach to constructing actions, this problem probably comes from subtleties involving the infinite number of fields. In the second approach, the problem probably comes from trying to define the correct phase space for the $a(x)$ - field [43]. In any case, this quantization problem is relevant for the actions proposed in this paper and certainly deserves further study.

Acknowledgements. The authors would like to thank K. Lechner, P. Pasti, S. Randjbar-Daemi, and M. Tonin for fruitful discussions, H. Nishino for pointing out an error in the supersymmetry transformations, E. Witten for pointing out difficulties with quantizing self-dual actions, S. de Alwis for comments, and P. Townsend for suggesting this work. I.B. and D.S. acknowledge partial support from grants of the Ministry of Science and Technology of Ukraine and the INTAS Grants N 93-493-ext, N 94-2317 and N 96-0308. N.B. acknowledges partial support from CNPq grant number 300256/94.

7. Appendix: SUSY transformation of $A^{(6)}$

To justify that the supersymmetry variation of $A^{(6)}$ has the right form, let us extend the field strengths (2.3) and (2.6) to a curved $D = 11$ superspace $Z^M = (x^{\underline{m}}, \Theta^\mu)$ where, to describe supergeometry, one introduces supervielbeins $E^A(Z) = dZ^M E_M^A = (E^{\underline{a}}, E^\alpha)$. Note that $E^{\underline{a}}|_{\Theta=0} = e^{\underline{a}}$ and $E^\alpha|_{\Theta=0} = \Psi^\alpha$.

The analysis of Bianchi identities [10], [44] shows that superfield strengths, which at $\Theta = 0$ reduce to (2.3) and (2.6), have only vector supervielbein components

$$\tilde{F}^{(4)} = \frac{1}{4!} E^{\underline{a}} E^{\underline{b}} E^{\underline{c}} E^{\underline{d}} \tilde{F}_{\underline{dcba}}^{(4)} \equiv dA^{(3)} - \frac{1}{2} E^{\underline{a}} E^{\underline{b}} E^\alpha E^\beta (\Gamma_{\underline{ab}})_{\alpha\beta}, \quad (7.1)$$

$$\begin{aligned} \tilde{F}^{(7)} &= \frac{1}{7!} E^{\underline{a}_1} \dots E^{\underline{a}_7} \tilde{F}_{\underline{a}_7 \dots \underline{a}_1}^{(7)} \\ &\equiv dA^{(6)} - A^{(3)} \wedge dA^{(3)} - \frac{2}{5!} E^{\underline{a}_1} \dots E^{\underline{a}_5} E^\alpha E^\beta (\Gamma_{\underline{a}_1 \dots \underline{a}_5})_{\alpha\beta}, \end{aligned} \quad (7.2)$$

where the last terms in (7.1) and (7.2) replace, respectively, $C^{(4)}$ and $C^{(7)}$ of (2.3) and (2.6) with their superform counterparts.

Now, by definition, a p -rank superform $W^{(p)}$ varies under supersymmetry transformations as follows:

$$\delta_\epsilon W^{(p)} = i_\epsilon dW^{(p)} + d(i_\epsilon W^{(p)}), \quad (7.3)$$

where i_ϵ defines the contraction of the spinor supervielbein components of the superform with the supersymmetry parameter ϵ^α . Hence, because of the constraints (7.1) and (7.2), the second term in (7.3) is absent from the supersymmetry variations of $\tilde{F}^{(4)}$ and $\tilde{F}^{(7)}$ (this just implies that they are supercovariant). Using these properties and substituting superfield analogues of the transformations (2.2) and (2.5) into the right hand side of the supersymmetry variation of (7.2), one can convince oneself that they correctly reproduce the supersymmetry variation (7.3) of the left hand side of (7.2).

References

- [1] E. Cremmer, B. Julia and J. Scherk, *Phys. Lett.* **B76** (1978) 409.
- [2] M. J. Duff and K. S. Stelle, *Phys. Lett.* **253B** (1991) 113.
- [3] R. Güven, *Phys. Lett.* **276B** (1992) 49.
- [4] O. Aharony, *Nucl. Phys.* **B476** (1996) 470 ;
E. Bergshoeff, M. de Roo and T. Ortín, *Phys. Lett.* **386B** (1996) 85.
- [5] P. Pasti, D. Sorokin and M. Tonin, *Phys. Lett.* **398B** (1997) 41.
- [6] I. Bandos, K. Lechner, A. Nurmagambetov, P. Pasti, D. Sorokin and M. Tonin, *Phys. Rev. Lett.* **78** (1997) 4332.
- [7] M. Aganagic, J. Park, C. Popescu and J. H. Schwarz, *Nucl. Phys.* **B496** (1997) 191.
- [8] D. Sorokin and P. K. Townsend, M-theory superalgebra from the M-5-brane, hep-th/9708003.
- [9] H. Nicolai, P. K. Townsend and P. van Nieuwenhuizen, *Lett. Nuov. Cim.* **30** (1981) 315.
- [10] R. D’Auria and P. Frè, *Nucl. Phys.* **B201** (1982) 101.
- [11] P.K. Townsend, *p-brane democracy*, hep-th/9507048.
- [12] S. P. de Alwis, Coupling of branes and normalization of effective actions in string/M-Theory, hep-th/9705139.
- [13] M. Perry and J. H. Schwarz, *Nucl. Phys.* **498** (1997) 47 ;
J. H. Schwarz, *Phys. Lett.* **B395** (1997) 191.
- [14] P. A. M. Dirac, *Proc. R. Soc.* **A133** (1931) 60, *Phys. Rev.* **74** (1948) 817 ;
N. Cabibbo and E. Ferrari, *Nuov. Cim.* **23** (1962) 1647 ;
J. Schwinger, *Phys. Rev.* **D3** (1971) 880.
- [15] D. Zwanziger, *Phys. Rev.* **D3** (1971) 880.
- [16] S. Deser, A. Gomberoff, M. Henneaux and C. Teitelboim, *Phys. Lett.* **B400** (1997) 80, hep-th/9702184.
- [17] N. Berkovits, *Phys. Lett.* **B395** (1997) 28, hep-th/9610134.
- [18] N. Berkovits and R. Medina, “Pasti-Sorokin-Tonin Actions in the Presence of Sources”, hep-th 9704093, to appear in *Phys. Rev. D*.
- [19] B. McClain, Y.S. Wu, and F. Yu, *Nucl. Phys.* **B343** (1990) 689, C. Wotzasek, *Phys. Rev. Lett.* **66** (1991) 129.
- [20] I. Martin and A. Restuccia, *Phys. Lett.* **B323** (1994) 311 ;
F. P. Devecchi and M. Henneaux, *Phys. Rev.* **D45** (1996) 1606.
- [21] N. Berkovits, *Phys. Lett.* **B388** (1996) 743, hep-th/9607070.
- [22] I. Bengtsson and A. Kleppe, *Int. J. Mod. Phys.* **A12** (1997) 3397.
- [23] P. Pasti, D. Sorokin and M. Tonin, *Phys. Rev.* **D52** (1995) 4277.
- [24] P. Pasti, D. Sorokin and M. Tonin, *Phys. Rev.* **D55** (1997) 6292.

- [25] R. Floreanini and R. Jackiw, *Phys. Rev. Lett.* **59** (1987) 1873 ;
M. Henneaux and C. Teitelboim, in Proc. Quantum mechanics of fundamental systems 2, Santiago, 1987, p. 79; *Phys. Lett.* **B206** (1988) 650.
- [26] J. H. Schwarz and A. Sen, *Nucl. Phys.* B411 (1994) 35.
- [27] M.J. Duff, B.E.W. Nilsson and C.N. Pope, *Phys. Rep.* 130 (1986) 1.
- [28] N. Berkovits and C. Hull, to appear ;
N. Nekrasov, private communication.
- [29] G. Dall'Agata, K. Lechner and M. Tonin, Covariant actions for N=1, D=6 supergravity theories with chiral bosons, hep-th/9710127.
- [30] P. Pasti, D. Sorokin and M. Tonin, Space-time symmetries in duality symmetric models. In Leuven Notes in Mathematical and Theoretical Physics, (Leuven University Press) Series B **V6** (1996) pp. 167–176, hep-th/9509053.
- [31] E. Bergshoeff, E. Sezgin and P. K. Townsend, *Phys. Lett.* **189B** (1987) 75; *Ann. Phys. (N.Y.)* **185** (1988) 330.
- [32] M. Blagojević and P. Senjanović, *Phys. Rep.* **157** (1988) 233.
- [33] I. Bengtsson, *Int. J. Mod. Phys.* **A12** (1997) 4869.
- [34] P.S. Howe and E. Sezgin *Phys. Lett.* **B394** (1997) 62;
P.S. Howe, E. Sezgin and P. West, *Phys. Lett.* **B399** (1997) 49 ;
C. S. Chu and E. Sezgin, M-fivebrane from the Open Supermembrane, hep-th/9710223.
- [35] I. Bandos, K. Lechner, A. Nurmagambetov, P. Pasti, D. Sorokin and M. Tonin, *Phys.Lett.* **B408** (1997) 135.
- [36] E. Witten, Five-brane effective action in M-theory, hep-th/9610234.
- [37] L. Bonora, C. S. Chu and M. Rinaldi, Perturbative anomalies of the M-5-brane, hep-th/9710063.
- [38] G. Dall'Agata, K. Lechner and D. Sorokin, Covariant Actions for the Bosonic Sector of D=10 IIB Supergravity, hep-th/9707044.
- [39] Y.-K. E. Cheung and Z. Yin, Anomalies, Branes and Currents, hep-th/9710206.
- [40] T. Kadoyoshi, S. Nojiri, S. D. Odintsov and A. Sugamoto, Vacuum polarization of supersymmetric D-brane in the constant electromagnetic field, hep-th/9710010.
- [41] A. Tseytlin, *Nucl. Phys.* **B469** (1996) 51.
- [42] D. Berman, SL(2,Z) duality of Born-Infeld theory from non-linear self-dual electrodynamics in 6 dimensions, hep-th/9706208.
- [43] E. Witten, private communication.
- [44] A. Candiello and K. Lechner, *Nucl. Phys.* **B412** (1994) 479.